









COVID-19 Isolation, Cleaning and Disinfection 2/4/2021

	<p>Airborne + Contact + Standard Precautions</p> <ul style="list-style-type: none"> Place mask on patient all the time if tolerated. Respirator (N95 or equivalent) Eye Protection: such as goggles or face shield Disposable gowns and disposable gloves Negative Pressure Room prioritized for aerosol generating procedures 	<p>Personal Protective Equipment (PPE) Disposal</p> <ul style="list-style-type: none"> All disposable PPE, other than N95, are discarded before leaving room. Only N95 mask is removed in the anteroom N95s that have been covered completely by a surgical mask or a face-shield can be considered for re-use (in which case they should be stored in the anteroom in paper bag labeled with staff name) Discard body fluid contaminated items as biohazard
	<p>Hand Hygiene</p> <ul style="list-style-type: none"> Alcohol-Based Hand Rub for at least 15 Seconds before and after all patient contact, contact with potentially infectious material, and before putting on and upon removal of PPE, including gloves 	<ul style="list-style-type: none"> If hands are visibly soiled, use soap and water before returning to alcohol-based hand rub (ABHR) 
	<p>Clorox Bleach Germicidal Wipes <u>3 Minutes of contact time</u> Sodium hypochlorite.....0.55%</p> <p>Micro-Kill Bleach Germicidal Wipes <u>3 Minutes of contact time</u> Sodium hypochlorite.....0.65%</p>	<ul style="list-style-type: none"> Hypochlorite is the disinfectant of choice for environmental decontamination for COVID-19 Patients or Suspects. All non-disposable equipment, including portable machines (portable X-ray, glucometers) and eye wear, must be disinfected for the required contact time
	<p>Hydrogen Peroxide Wipes: <u>5 Minutes of contact time</u> Hydrogen Peroxide.....1.4%</p>	<ul style="list-style-type: none"> Hydrogen Peroxide is the disinfectant of choice for environmental decontamination for COVID-19 Patients or Suspects in OR.
	<p>Super Sani-Cloth WIPES <u>2 Minutes</u> Isopropyle Alcohol.....55% dimethyl benzyl ammonium chlorides 0.5%</p>	<ul style="list-style-type: none"> General purpose disinfection of hard nonporous surfaces and equipment made of plastic, Formica laminate and glass Disinfection of Nova StatStrip Glucometers
	<p>Sani-Cloth AF3 <u>3 Minutes</u> Dimethyl ethylbenzyl ammonium chlorides.....0.14% Dimethyl benzyl ammonium chlorides.....0.14%</p>	<p>Disinfection of hard, nonporous surfaces and equipment when the manufactures require alcohol-free disinfectant such as for Ultrasound machine, user preference is allowed for the lack of smell and for rubber surfaces.</p>
	<p>Room Entry After Patient Discharge</p> <ul style="list-style-type: none"> Use PPE while cleaning the room after discharge 	<ul style="list-style-type: none"> 10 minutes if no aerosol-generating procedure (AGP) 45 Minutes for AIIR used for aerosol-generating procedure (10-12 air exchanges per hour) 3.5 hours for regular patient room in VCMC used for aerosol-generating procedure (2 air exchanges per hour) 1.5 hours for regular patient room used for aerosol-generating procedure in SPH (6 air exchanges per hour)